



...go modern



use

PRIMACAINE

a brand of METABUTOXYCAINE HCl



Primacaine HCl (Metabutoxycaine HCl) — the modern local anesthetic — offers the qualities you look for in a local anesthetic ...*very rapid anesthesia...great potency...adequate margin of safety...ideal duration...and excellent tolerance by all types of patients.* (Available in 2 concentrations of epinephrine — epinephrine 1:60,000 for general use; and epinephrine 1:125,000 for special cases.)

If you do not use PRIMACAINE HCl routinely, we suggest that you try it and compare in your day-by-day practice. Clinical data on request.

Novocol

CHEMICAL MFG. CO., Inc.—BROOKLYN 7, NEW YORK

Canadian Laboratories, TORONTO 5, ONTARIO

*Primacaine HCl is the registered trade mark of Novocol Chem. Mfg. Co., Inc., designating its product, 2'-diethylamino-ethyl-2-butoxy-3 amino benzoate HCl.

earlier total recovery **Neraval** SODIUM

IN INTRAVENOUS ANESTHESIA FOR ORAL SURGERY

*Confirmed by
recent clinical
observations*

"Our initial impression [with NERAVAL] over a series of 200 cases for prolonged one to two hour oral rehabilitation cases, has impressed us with the quicker recovery time, less hangover sensation and earlier ambulation."

Spiro, S. R.: Personal Communication.

Based on administration to over 1000 patients aged 3 months to 89 years for all types of surgery, including oral surgery, one investigator stated:

"When used as an induction agent with the semiclosed cyclopropane-nitrous oxide-oxygen system (The Balanced Mixture) Neraval is without a doubt the best ultrashort-acting barbiturate available today. Patients not only awoken more quickly, but they are rationally and coherently awake more quickly than that seen with any other barbiturate used as the induction agent."

Shane, S. M.: Personal Communication.

"Neraval Sodium was found to be markedly more short-acting than thiopental sodium in a series of 427 ambulatory dental patients.

"...The prolonged narcosis frequently experienced after thiopental was not noted with Neraval.

"...The percentage of total recoveries within one hour in the Neraval group was almost twice that of the thiopental group."

Krasner, H. A.: Personal Communication.

"The prolonged depression, which frequently follows thiobarbiturate administration, was not observed with Neraval.

"Many patients who had previously experienced post-anesthetic side effects with thiopental anesthesia comment on the *freedom from* discomfort and drowsiness which characterized recovery from Neraval.

"...The reduction in patient hours which occurs with Neraval permits increased scheduling and lessens the load of postanesthetic nursing supervision."

Gallagher, C. J.: Recovery after a new intravenous anesthesia "Neraval," (methitural sodium), in 109 dental patients, J. New Jersey Dent. Soc. 28:7 (July) 1957.

Supplied

NERAVAL Sodium Sterile Powder—vials of 1 Gm. and 2 Gm., boxes of 6 and 25; vial of 5 Gm., boxes of 1 and 25.

Literature will be sent on request.

NERAVAL® Sodium, brand of methitural sodium.

SCHERING CORPORATION • BLOOMFIELD, NEW JERSEY

Schering

NE-J-2117

New A.D.S.A. Members**(continued)**

Dr. Rees G. Russell
Bristol Memorial Building
Bristol, Tennessee

Dr. Joseph H. Rutherford
2223 Austin Avenue
Waco, Texas

Dr. Donald N. Shapiro
201 Macole Building
22720 Woodward Avenue
Ferndale, Michigan

Dr. Thomas Edgar Sikes, Sr.
916 Southeastern Building
Greensboro, North Carolina

Col. William B. Simms
Tokyo, U.S. Army Hospital
A.P.O. 500
San Francisco, California

Dr. Roy Galen Slack
4747 Mission Boulevard
Suite 3
San Diego 9, California

Dr. John R. Swanson
923 Medical Arts
Dallas 1, Texas

Dr. Lucian Szmyd
School of Aviation Medicine
U.S.A.F.
Randolph A.F.B., Texas

Dr. Mack Taylor
103 W. Harrison Street
Danville, Illinois


Dr. Kenneth W. Thomas
1015 Shook at Hildebrand
San Antonio, Texas

Capt. A. S. Turville
U. S. Naval Hospital
Oakland 14, California

Dr. Robert F. White
932 Maple Avenue
Franklin Grove, Illinois

Dr. Harold Zubrow
3347 Forbes Street
Pittsburgh 13, Pennsylvania

The SIMMONDS ARM BOARD



for INTRAVENOUS ANESTHESIA

The Simmonds Arm Board* is an excellent apparatus which facilitates the use of intravenous anesthetic agents† at the dental chair or the operating table. As portrayed in the above photograph, a 10 cc. syringe is utilized for the injection of the agent. Withdrawal of the plunger automatically fills the syringe from the reservoir which hangs immediately below the arm board. This reservoir is inconspicuous and out of the operating field.

†Type and percentage of solution can be varied.

* The Simmonds Arm Board is a mechanical aid to assist with the administration of intravenous agents in general anesthesia.

Cost	Includes	\$65.00
	Syringes	
	Valves	
	Venotubes	
	Venopaks	

Manufactured by
DAVID SIMMONDS
Instrument Maker

17 West 60th St. - New York 23, N.Y.

*For the dentist who
wants to please...*



RAVOCAINE^{HCl} AND **NOVOCAIN**^{HCl}
Brand of propoxycaine HCl Brand of procaine HCl
WITH **NEO-COBEFRIN**
Brand of levo-nordefrin

Extremely Rapid Onset - Unsurpassed Depth - Moderate Duration

***Plus* A New High in Patient Tolerance**

The extremely rapid onset; unsurpassed depth and moderate duration, which have made the other Ravocaine-Novocain combinations so famous, also distinguish this solution. Use of the better tolerated vaso-constrictor, NEO-COBEFRIN, the levorotatory form of time-tested Cobefrin,* produces the new, higher level of patient comfort and tolerance.

Order your supply of Ravocaine HCl 0.4% and Novocain 2% with Neo-Cobefrin 1:20,000 today. . . it's ideal for restorative and short surgical procedures.

COOK-WAITE
Laboratories, Inc.

1450 BROADWAY, NEW YORK 18, N. Y.

* Cobefrin (brand of nordefrin), Neo-Cobefrin, Novocain and Ravocaine are the trademarks (Reg. U.S. Pat. Off.) of Sterling Drug Inc.